

Short report

Open Access

A confusing case of canine vector-borne disease: clinical signs and progression in a dog co-infected with *Ehrlichia canis* and *Bartonella vinsonii ssp. berkhoffii*

Edward B Breitschwerdt*, Ricardo G Maggi

Address: Department of Clinical Sciences, College of Veterinary Medicine, North Carolina State University, 4700 Hillsborough Street, Raleigh, NC 27606, USA

Email: Edward B Breitschwerdt* - ed_breitschwerdt@ncsu.edu

*Corresponding Author

from 4th International Canine Vector-Borne Disease Symposium
Seville, Spain. 26–28 March 2009

Published: 26 March 2009

Parasites & Vectors 2009, **2**(Suppl 1):S3 doi:10.1186/1756-3305-2-S1-S3

This article is available from: <http://www.parasitesandvectors.com/content/2/S1/S3>

© 2009 Breitschwerdt and Maggi; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Bartonella spp. are important pathogens in human and veterinary medicine, and bartonellosis is considered as an emerging zoonosis that is being reported with increasing frequency. Of 22 known species and subspecies of *Bartonella*, seven have been isolated from dogs, causing disease manifestations similar to those seen in human beings. The wide variety of clinical signs and the possible chronic progression of disease manifestations are illustrated in the case of an infected Labrador retriever. Here, the authors discuss the seemingly diverse spectrum of disease manifestations, the co-infections of *Bartonella* spp. with other vector-borne pathogens (mainly *Ehrlichia* spp. or *Babesia* spp.) and the difficulties in microbiological confirmation of an active *Bartonella* infection, all of which make the disease pathogenesis and clinical diagnosis more problematic.

Findings

Historically, the most prevalent *Bartonella* species in dogs, *B. vinsonii ssp. berkhoffii*, was initially isolated from a dog with endocarditis and intermittent epistaxis in 1993 [1,2]. Subsequently, this pathogen has also been associated with cardiac arrhythmias, myocarditis, granulomatous rhinitis, anterior uveitis and chorioiditis [1,3-6]. Other *Bartonella* species which have also been associated with pathology and clinical signs in dogs, including endocarditis, hepatic disease and sudden death are: *B. henselae* [7-11],

B. clarridgeiae [9,12], *B. washoensis* [13], *B. elizabethae* [14] and *B. quintana* [15]. The first and the last, *B. henselae* and *B. quintana*, were also detected in the blood or lymph nodes of healthy dogs and dogs suffering from lymphoma [8]. More recent studies indicate that *B. henselae* may be the most frequent cause of bartonellosis in dogs.

The numerous disease manifestations in dogs, many of which are similar to those seen in humans, are listed in Table 1.

Table 1 - *Bartonella* infection in dogs.

Species/subspecies	Associated manifestations of infection [23-31]
<i>Bartonella vinsonii</i> subsp. <i>berkhoffii</i>	Endocarditis* Cardiac arrhythmias* Myocarditis* Polyarthritis* Granulomatous rhinitis Anterior uveitis Chorioretinitis* Meningoencephalitis* Anaemia/Thrombocytopenia*
<i>Bartonella henselae</i>	Peliosis hepatitis* Granulomatous hepatitis* Generalised pyogranulomatous lymphadenitis* Panniculitis Endocarditis* Polyarthritis* Idiopathic effusions
Other <i>Bartonella</i> species	Endocarditis*
<i>Bartonella clarridgeiae</i>	Hepatic disease
<i>B. washoensis</i>	Weight loss
<i>B. elizabethae</i>	
<i>B. quintana</i>	

* Denotes disease manifestations reported in dogs and human beings.

Bartonella spp. are transmitted by bites and scratches of infected animals to other hosts. In addition, arthropod vectors may play the most important role in the transmission of these organisms. *B. henselae*, the agent of cat-scratch disease (CSD), has been isolated from cat fleas, and transmission by ticks has been proposed [16,17]. Although tick transmission of *B. vinsonii* ssp. *berkhoffii* has been suspected for over a decade based upon epidemiological evidence, this has not been proven [17]. Evidence suggests that dogs, coyotes and gray foxes may be the reservoir hosts for *B. vinsonii* ssp. *berkhoffii* [18,19]. Due to the mode of transmission a co-infection with other arthropod-borne pathogens is possible and may mask or alter the typical clinical signs of each of these pathogens. The following case of a chronically ill Labrador retriever is shortly outlined to illustrate the potential confusion and clinical challenges induced by tick-borne infections:

A 3-year-old, spayed, female Labrador retriever was referred to the North Carolina State University Veterinary Teaching Hospital for evaluation of a protracted illness that had begun approximately 9 months earlier. Initially, the dog was lethargic and intermittently inappetent and had a shifting-leg lameness suggestive of polyarthritis. On the initial examination prior to referral, the dog was mildly anaemic (haematocrit, 36%) and hyperprotein-aemic (8.0 g/dl), but showed no obvious physical exami-

nation abnormalities. Two weeks later, the dog developed a grand mal seizure, accompanied by spontaneous urination. During the subsequent month, lethargy continued and weight loss was recorded (5.0 kg). On a second veterinary examination numerous intradermal haemorrhages on the neck and trunk were noticed as well as anaemia (haematocrit: 32%) with spherocytosis, mild thrombocytopenia (platelets: 180,000/ μ l), hyperglobulin-aemia (serum globulin concentration: 4.7 g/dl), proteinuria, and hemoglobinuria (urine specific gravity: 1.034; 4+ protein; 4+ blood; 4 to 8 erythrocytes per high-power field) and physiological leukocyte counts and differential cell numbers. Seroreactivity to *Ehrlichia canis*, *Rickettsia rickettsii* and nuclear (antinuclear antibodies) antigens by indirect fluorescent-antibody assays (reciprocal titers, 50, 256 and 640, respectively) were detected. Ehrlichiosis and systemic lupus erythematosus were diagnosed by the referring veterinarian.

Treatment was initiated with tetracycline hydrochloride (750 mg 3 times daily for 14 days) and prednisone (40 mg twice daily (BID) for 3 days, then gradually tapered to 15 mg every other day).

During the next 3 months, episodes of listlessness and epistaxis occurred and on referral examination the dog was lethargic with slight tachycardia, a hyperdynamic arterial pulse accompanied by arterial pulse deficits and occasional premature beats. A diastolic and a systolic heart murmur were recorded. On electrocardiogram and spectral Doppler echocardiography extrasystoles and severe aortic as well as mild mitral valve insufficiency were diagnosed, already resulting an increase in interstitial and alveolar pulmonary infiltrates in the accessory lung lobe (thoracic radiographs). Laboratory parameters proved anaemia (haematocrit: 33%; mild anisocytosis and macrocytosis), thrombocytopenia (platelet count: 121,000/ μ l), leukocyte counts at 12,300/ μ l, with normal differential cell numbers, hypoalbumin-aemia (serum albumin: 2.6 g/dl), azotaemia (blood urea nitrogen: 34 mg/dl) and hypokalaemia (serum potassium: 3.9 g/dl). Urinalysis again revealed proteinuria and haemoglobinuria (specific gravity: 1.011; 3+ protein; 2+ blood), haematuria (5 to 10 erythrocytes per high-power field), pyuria (5 to 10 leukocytes per high-power field) and bacteriuria. Aerobic and anaerobic blood cultures during a 24-h period as well as terminal subcultures and Gram stains performed after 7 days failed to grow or identify bacteria. Blood cultured simultaneously by the lysis centrifugation technique grew a fastidious, gram-negative organism. Seroreactivity (indirect fluorescent-antibody assay) to *E. canis* and *R. rickettsii* was again positive (reciprocal titers, 64 and 128, respectively) and a specific antibody response to *E. canis* was confirmed by Western immunoblot analysis.

Treatment for bacterial endocarditis accompanied by congestive heart failure consisted of enrofloxacin (306 mg BID), doxycycline (400 mg BID), clavulanate-potentiated amoxicillin (530 mg BID), furosemide (40 mg BID), digoxin (0.25 mg BID) and enalapril (20 mg BID). But because of a lack of substantial clinical improvement, intractable epistaxis and the poor long-term prognosis associated with vegetative valvular endocarditis, the owners elected euthanasia 17 days following discharge.

A lysis centrifugation technique finally isolated *B. vinsonii* ssp. *berkhoffii* genotype I in this dog by blood culture so that it was suggested that the dog was co-infected with *E. canis* and *B. vinsonii* ssp. *berkhoffii*. Reactivity to *Rickettsia* antigens is suggested to be cross-reactive with *Bartonella* antigens and anti-nuclear antibodies can occur with an increased frequency in dogs that are seroreactive to *B. vinsonii* ssp. *berkhoffii* and *E. canis* antigens [20].

The above-described case report highlights three problems related to infections with *Bartonella* spp.: (1) A large variety of disease manifestations makes a clinical diagnosis more difficult (diagnostic indications that support the testing for *Bartonella* infection are listed in Table 2); (2) co-infections of *Bartonella* spp. with other vector-borne pathogens particularly *Ehrlichia* spp. or *Babesia* spp. may influence the pathogenesis of those diseases and alternatively co-infection may also alter clinical signs of bartonellosis, creating substantial challenges for the clinician in regard to accurate diagnosis and directed medical management; (3) microbiological confirmation of an active *Bartonella* infection is difficult. Concerning the last point, and as is seen with other intracellular pathogens that induce chronic infection after vector-borne transmission, this aspect of diagnostic confirmation can be very challenging [7]. Ideally, diagnosis should be confirmed by culturing the organism from biopsy specimens, blood or fluid samples that are aseptically obtained. Due to the fastidious growth characteristics of *Bartonella* spp. it is important to avoid contamination of diagnostic specimens with skin flora or other rapidly growing bacteria. Unlike cats, which may easily have a bacterial load of 100,000 copies/ μ l, dogs are likely to have levels that are 100- to 1000-fold lower [7]. The presence of antibodies can only be used to infer prior exposure, but a substantial number of infected dogs do not have detectable antibodies. PCR amplification of *Bartonella* DNA following direct extraction from patient samples is also relatively insensitive. Two media have been developed to overcome the problem of slow-growing pathogens and unavailable liquid medium: "BAPGM" medium [21] and another medium developed by Riess *et al.* [22]. Nevertheless, culturing remains very time-consuming and when the need for a diagnosis is urgent, treatment should be started after obtaining appropriate

Table 2 - Differential diagnostic indications that support consideration of testing for *Bartonella* infection.

Granulomatous inflammatory lesions
Unexplained reactive lymphadenopathy
Endocarditis
Myocarditis
Polyarthritis
Immune-mediated haemolytic anaemia
Immune-mediated thrombocytopenia
Eosinophilia
Splenomegaly
Epistaxis
Idiopathic cavity effusions
Unexplained neurologic disease
Fever of unknown origin
Vasculitis
Chronic hepatitis

diagnostic specimens. Serum antibodies can be detected in both healthy and clinically ill dogs and thus do not always correlate with illness. Therefore antibodies may only reflect prior exposure, but do not necessarily document that *Bartonella* is the cause of the current disease.

Bartonella spp. infections in dogs are increasingly reported. Dogs show a variety of clinical manifestations ranging from asymptomatic to severe disease, a fact that makes clinical diagnosis quite difficult. Studies have shown that infection with *B. vinsonii* ssp. *berkhoffii* can induce polyarthritis, seizures, vasculitis, epistaxis and endocarditis in dogs (all of which occurred in progression in the illustrated case report).

Other underlying co-infections which are also vector-borne, such as infections with *Ehrlichia canis* and *Babesia* spp., may further influence the clinical presentation and also influence the prognosis, particularly if all co-infected organisms are not detected or specifically treated. Finally, laboratory diagnosis of *Bartonella* infection in dogs can also be quite difficult and time-consuming (if performed by culturing). Many of these aspects could be illustrated in this case report of a confusing tick-borne disease and should be kept in mind by veterinary clinicians.

Competing interests

Drs Breitschwerdt and Maggi are scientific officers for Galaxy Diagnostics, a newly formed company devoted to enhanced detection of *Bartonella* spp. infection in animals and human beings and located in Raleigh, NC, USA.

Authors' contributions

Drs Breitschwerdt and Maggi have collaborated for the past five years on studies to enhance the diagnosis, treatment and prevention of *Bartonella* infection in animals and human beings. Each contributed equally to the content of this case report.

Acknowledgements

We thank the members of the Vector-Borne Diseases Diagnostic Laboratory and the Intracellular Pathogens Research Laboratory for providing the diagnostic and research expertise that facilitated the evaluation of this dog.

This article is published as part of *Parasites & Vectors* Volume 2 Supplement 1, 2009: Proceedings of the 4th International Canine Vector-Borne Disease Symposium. The full contents of the supplement are available online at <http://www.parasitesandvectors.com/supplements/2/S1>.

Publication of the supplement has been sponsored by Bayer Animal Health GmbH.

References

- Breitschwerdt EB, Kordick DL, Malarkey DE, Keene B, Hadfield TL, Wilson K: **Endocarditis in a dog due to infection with a novel *Bartonella* subspecies.** *J Clin Microbiol* 1995, **33**:154-160.
- Kordick DL, Swaminathan B, Greene CT, Wilson KH, O'Connor S, Hollis DG, Matar GM, Malcolm GB, Hayes PS, Hadfield TL, Breitschwerdt EB, Brenner DJ: ***Bartonella vinsonii* subsp. *berkhoffii* subsp. nov., isolated from dogs; *Bartonella vinsonii* subsp. *vinsonii*; and emended description of *Bartonella vinsonii*.** *Int J Syst Bacteriol* 1996, **46**:704-709.
- Breitschwerdt EB, Atkins CE, Brown TT, Kordick DL, Snyder PS: ***Bartonella vinsonii* subsp. *berkhoffii* and related members of the alpha subdivision of the Proteobacteria in dogs with cardiac arrhythmias, endocarditis, or myocarditis.** *J Clin Microbiol* 1999, **37**:3618-3626.
- Michau TM, Breitschwerdt EB, Gilger BC, Davidson MG: ***Bartonella vinsonii* subspecies *berkhoffii* as a possible cause of anterior uveitis and choroiditis in a dog.** *Vet Ophthalmol* 2003, **6**:299-304.
- Pappalardo BL, Brown T, Gookin JL, Morrill CL, Breitschwerdt EB: **Granulomatous disease associated with *Bartonella* infection in 2 dogs.** *J Vet Intern Med* 2000, **14**:37-42.
- Pappalardo BL, Brown TT, Tompkins M, Breitschwerdt EB: **Immunopathology of *Bartonella vinsonii* (*berkhoffii*) in experimentally infected dogs.** *Vet Immunol Immunopathol* 2001, **83**:125-147.
- Duncan AV, Maggi RG, Breitschwerdt EB: **A combined approach for the enhanced detection and isolation of *Bartonella* species in dog blood samples: pre-enrichment liquid culture followed by PCR and subculture onto agar plates.** *J Microbiol Methods* 2007, **69**:273-281.
- Duncan AV, Marr HS, Birkenheuer AJ, Maggi RG, Williams LE, Correa MT, Breitschwerdt EB: ***Bartonella* DNA in the blood and lymph nodes of Golden Retrievers with lymphoma and in lymph controls.** *J Vet Intern Med* 2008, **22**:89-95.
- Gillespie TN, Washabau RJ, Goldschmidt MH, Cullen JM, Rogala AR, Breitschwerdt EB: **Detection of *Bartonella henselae* and *Bartonella clarridgeiae* DNA in hepatic specimens from two dogs with hepatic disease.** *J Am Vet Med Assoc* 2003, **222**:47-51, 35.
- Kitchell BE, Fan TM, Kordick D, Breitschwerdt EB, Wollenberg G, Lichtensteiger CA: **Peliosis hepatis in a dog infected with *Bartonella henselae*.** *J Am Vet Med Assoc* 2000; **216**:519-523, 517.
- Saunders GK, Monroe VE: **Systemic granulomatous disease and sialometaplasia in a dog with *Bartonella* infection.** *Vet Pathol* 2006, **43**:391-392.
- Chomel BB, Mac Donald KA, Kasten RW, Chang CC, Wey AC, Foley JE, Thomas WP, Kittleson MD: **Aortic valve endocarditis in a dog due to *Bartonella clarridgeiae*.** *J Clin Microbiol* 2001, **39**:3548-3554.
- Chomel BB, Wey AC, Kasten RW: **Isolation of *Bartonella washoensis* from a dog with mitral valve endocarditis.** *J Clin Microbiol* 2003, **41**:5327-5332.
- Boulouis HJ, Chang CC, Henn JB, Kasten RW, Chomel BB: **Factors associated with the rapid emergence of zoonotic *Bartonella* infections.** *Vet Res* 2005, **36**:383-410.
- Kelly P, Rolain JM, Maggi R, Sontakke S, Keene B, Hunter S, Lepidi H, Breitschwerdt KT, Breitschwerdt EB: ***Bartonella quintana* endocarditis in dogs.** *Emerg Infect Dis* 2006, **12**:1869-1872.
- Billeter SA, Levy MG, Chomel BB, Breitschwerdt EB: **Vector transmission of *Bartonella* species with emphasis on the potential for tick transmission.** *Med Vet Entomol* 2008, **22**:1-15.
- Cotté V, Bonnet S, Le Rhun D, Le Naour E, Chauvin A, Boulouis HJ, Lecuelle B, Lilin T, Vayssier-Taussat M: **Transmission of *Bartonella henselae* by *Ixodes ricinus*.** *Emerg Infect Dis* 2008, **14**:1074-1080.
- Pappalardo BL, Correa MT, York CC, Peat CY, Breitschwerdt EB: **Epidemiologic evaluation of the risk factors associated with exposure and seroreactivity to *Bartonella vinsonii* in dogs.** *Am J Vet Res* 1997, **58**:467-471. Risk factors for epidemiology.
- Chang C-C, Kasten RW, Chomel BB, Simpson DC, Hew CM, Kordick DL, Heller R, Piemont Y, Breitschwerdt EB: **Coyotes (*Canis latrans*) as the reservoir for a human pathogenic *Bartonella* sp.: molecular epidemiology of *Bartonella vinsonii* subsp. *berkhoffii* infection in coyotes from central coastal California.** *J Clin Microbiol* 2000, **38**:4193-4200.
- Maggi RG, Chomel B, Hegarty BC, Henn J, Breitschwerdt EB: **A *Bartonella vinsonii berkhoffii* typing scheme based upon 16S-23S ITS and Pap31 sequences from dog, coyote, gray fox, and human isolates.** *Mol Cell Probes* 2006, **20**:128-134.
- Smith BE, Tompkins MB, Breitschwerdt EB: **Antinuclear antibodies can be detected in dog sera reactive to *Bartonella vinsonii* subsp. *berkhoffii*, *Ehrlichia canis*, or *Leishmania infantum* antigens.** *J Vet Intern Med* 2004, **18**:47-51.
- Maggi RG, Duncan AV, Breitschwerdt EB: **Novel chemically modified liquid medium that will support the growth of seven *Bartonella* species.** *J Clin Microbiol* 2005, **43**:2651-2655.
- Riess T, Dietrich F, Schmidt KV, Kaiser PO, Schwarz H, Schäfer A, Kempf VA: **Analysis of a novel insect-cell culture medium-based growth medium for *Bartonella* species.** *Appl Environ Microbiol* 2008, **74**:5224-5227.
- Chomel BB, Boulouis HJ, Maruyama S, Breitschwerdt EB: ***Bartonella* spp. in pets and effect on human health.** *Emerg Infect Dis* 2006, **12**:389-394.
- Chomel BB, Kasten RW, Sykes JE, Boulouis HJ, Breitschwerdt EB: **Clinical impact of persistent *Bartonella* bacteremia in humans and animals.** *Ann NY Acad Sci* 2003, **990**:267-278.
- Mexas AM, Hancock SI, Breitschwerdt EB: ***Bartonella henselae* and *Bartonella elizabethae* as potential canine pathogens.** *J Clin Microbiol* 2002, **40**:4670-4674.
- Morales SC, Breitschwerdt EB, Washabau RJ, Matisse I, Maggi RG, Duncan AV: **Detection of *Bartonella henselae* DNA in two dogs with pyogranulomatous lymphadenitis.** *J Am Vet Med Assoc* 2007, **230**:681-685.
- Goodman RA, Breitschwerdt EB: **Clinicopathologic findings in dogs seroreactive to *Bartonella henselae* antigens.** *Am J Vet Res* 2005, **66**:2060-2064.
- Diniz PP, Maggi RG, Schwartz DS, Cadenas MB, Bradley JM, Hegarty B, Breitschwerdt EB: **Canine bartonellosis: serological and molecular prevalence in Brazil and evidence of co-infection with *Bartonella henselae* and *Bartonella vinsonii* subsp. *berkhoffii*.** *Vet Res* 2007, **38**:697-710.
- Mellor PJ, Fetz K, Maggi RG, Haugland S, Dunning M, Villiers EJ, Mellanby RJ, Williams D, Breitschwerdt E, Herrtage ME: **Alpha1-proteinase inhibitor deficiency and *Bartonella* infection in association with panniculitis, polyarthritis, and meningitis in a dog.** *J Vet Intern Med* 2006, **20**:1023-1028.
- Breitschwerdt EB, Blann KR, Stebbins ME, Muñana KR, Davidson MG, Jackson HA, Willard MD: **Clinicopathologic abnormalities and treatment response in 24 dogs seroreactive to *Bartonella vinsonii* (*berkhoffii*) antigens.** *J Am Anim Hosp Assoc* 2004, **40**:92-101.